

When are There Infinitely Many Irreducible Elements in a Principal Ideal Domain?

Fabrizio Zanello

It has been a well-known fact since Euclid's time that there exist infinitely many rational primes. Two natural questions arise: In which other rings, sufficiently similar to the integers, are there infinitely many irreducible elements? Is there a unifying algebraic concept that characterizes such rings?

The purpose of the present note is to place the fact concerning the infinity of primes into a more general context, one that also includes the interesting case of the factorial domains (unique factorization domains) of algebraic integers in a number field. We show that, if A is a Principal Ideal Domain (PID, for short), then the fact that A contains infinitely many (pairwise nonassociate) irreducible elements is equivalent to the property that every maximal ideal in the polynomial ring $A[x]$ has the same (maximal) height.

We begin by recalling some basic definitions. (We assume that all rings are commutative with an identity element, denoted by 1). The *Jacobson radical* $J(R)$ of a ring R is the intersection of all the maximal ideals of R , while the *nilradical* $\sqrt{0}$ of R is the intersection of all the prime ideals of R . The latter can also be described as the set of all nilpotent elements of R (see [1, Proposition 1.8, p. 5]). The

height of a prime ideal P in R is the supremum of the lengths of the chains $P_0 \subset P_1 \subset \dots \subset P_r = P$ of prime ideals of R that end at P . The *Krull-dimension* $\dim R$ of R is the supremum of the lengths of all the chains of prime ideals of R , or, equivalently, the supremum of the heights of all the prime ideals P in R . For instance, a field has Krull-dimension 0, while a PID A has Krull-dimension 1, since every prime ideal of A different from (0) has height 1. Finally, two elements a and b of R are *associates* if there exists a unit u in R such that $a = ub$. Otherwise, we say that a and b are *nonassociates*.

Lemma 1. *Let R be a domain. Then for the ring $R[x]$ it is the case that $J(R[x]) = \sqrt{0} = (0)$.*

Proof. Of course $\sqrt{0} = (0)$, for $R[x]$ contains no nonzero nilpotent elements. Now let $f(x)$ be a member of $J(R[x])$. Then $xf(x) + 1$ is a unit: otherwise $xf(x) + 1$ would belong to some maximal ideal M in $R[x]$, and since $f(x)$ lies in M , this would place 1 in M , a contradiction. Since $R[x]$ is a domain, this implies that $xf(x) + 1$ is a constant polynomial, so $f = 0$. $\square$

Actually, the conclusion of Lemma 1 can be extended to an arbitrary ring R . It is not difficult to show (but the argument is longer) that in every $R[x]$ we have $J(R[x]) = \sqrt{0}$ (see [1, Exercise 4, p. 11]).

Theorem 2. *Let A be a PID. Then the following statements are equivalent:*

(i) *every maximal ideal of $A[x]$ has height 2;*

(ii) A has infinitely many pairwise nonassociate irreducible elements.

Proof. Let M be a maximal ideal in $A[x]$ and set $P = M \cap A$. Then P is a prime ideal of A . Suppose for the moment that $P = (0)$. We claim that M has height 1 in this situation. Choose an element $g(x)$ in M of least degree. Then $\deg g(x) > 0$, and we can assume that $g(x)$ is irreducible, since M is a prime ideal.

Let K be the quotient field of A . If $f(x)$ belongs to M , then in $K[x]$ we can write

$$f(x) = g(x)q_1(x) + r_1(x),$$

where either $r_1(x) \equiv 0$ or $\deg r_1(x) < \deg g(x)$. Therefore we obtain

$$f(x) = g(x)\frac{q(x)}{a} + \frac{r(x)}{a},$$

or

$$af(x) = g(x)q(x) + r(x),$$

with $q(x)$ and $r(x)$ in $A[x]$ and a in $A - \{0\}$. This implies that $r(x)$ lies in M , whence $r(x) \equiv 0$, because $\deg r(x) \geq \deg g(x)$ by the choice of $g(x)$. Therefore $af(x) \in (g(x)) \subset M$. We infer that $f(x)$ belongs to $(g(x))$: $(g(x))$ is a prime ideal, so either a or $f(x)$ belongs to it, but prime ideals in $A[x]$ cannot contain nonzero constants. Hence $M = (g(x))$, from which it follows that M has height 1 in case $P = (0)$.

Proof of (i) $\Rightarrow$ (ii). Assume that every maximal ideal M in $A[x]$ has height 2. From what we have just shown, it follows that every such M must contain some nonzero constant, hence must include at least one irreducible element of A .

Suppose now that there are only finitely many pairwise nonassociate irreducible elements of A and that $p_1, p_2, \dots, p_s$ is a complete list of representatives of irreducible elements. Then $p_1 p_2 \cdots p_s$ belongs to every maximal ideal M in $A[x]$, that is, to $J(A[x])$. Because A is a domain, Lemma 1 asserts that $J(A[x]) = (0)$, a contradiction. This proves (ii).

Proof of (ii) $\Rightarrow$ (i). Let M again be an arbitrary maximal ideal of $A[x]$. We want to show that M has height 2.

If $P = M \cap A = (0)$, then we have already seen that $M = (g(x))$ for some irreducible $g(x)$ in $A[x]$. Since M is a maximal ideal, for every irreducible p in A we must have $(g(x), p) = (1)$ (i.e., $A[x]/(g(x), p) = 0$), and therefore $\overline{g(x)}$ must be a constant in $(A/(p))[x]$ for each such p . In other words, every coefficient of $g(x)$ other than the constant term is divisible by every irreducible element p of A . This is a contradiction, since we are assuming that there are infinitely many pairwise nonassociate irreducible elements of A and since, as a PID, A is a factorial domain.

Hence $P = M \cap A \neq (0)$, so $P = (p)$ for some irreducible element p in A . Moreover M is not principal: if it were, it would coincide with (p) , which is impossible because $A[x]/(p) \simeq (A/(p))[x]$ is not a field.

Hence M strictly contains the prime ideal (p) , and therefore its height is at least 2. But

$$\dim A[x] = \dim A + 1 = 2,$$

since A is Noetherian (see [1, Exercise 7, p. 126]). Thus the height

of M is exactly 2, as required by (i). The proof of the theorem is complete. $\square$

ACKNOWLEDGEMENTS. I wish to express my warm gratitude to Professor Ram Murty for an interesting discussion that led to the idea of writing this paper.

References

- [1] M.F. Atiyah and I.G. Macdonald, *Introduction to Commutative Algebra*, Addison-Wesley, Reading, MA, 1969.

Fabrizio Zanello

Department of Mathematics and Statistics, Queen's University

Kingston, Ontario

K7L 3N6 Canada

fabrizio@mast.queensu.ca